

Outline


- Structural and functional organization of chromosomes
- DNA packaging and structural organization
- Overview of kinetochore structure and functions

Outline

- Structural and functional organization of chromosomes
- DNA packaging and structural organization
- Overview of kinetochore structure and functions



The Problem

- Spacing between base pairs $\approx 3.4\text{\AA}$
- For human genome, approximately 3.2 billion base pairs
- Total length $\approx 3.4 \times 10^{-10} \times 3.2 \times 10^9 \times 2 \approx 2.2\text{m}$
- Diameter of a nucleus: $5 \sim 10 \times 10^{-6}\text{m}$
- Access to genetic information must be regulated.





Chromosome Organization in the Cell Cycle

- DNA are packaged with associated proteins into chromosomes.
- Chromosomes undergo dramatic reorganization when cells divide.




Chromosome Related Nomenclature (I)

- Chromatin: the complex of DNA, histones, and nonhistone proteins within the nucleus of a eukaryotic cell (DNA persistence length ~50nm). The material of which chromosomes are made.
- Chromatid: one of the two copies of a replicated chromosome that is joined at the centromere to the other copy. The two identical chromatids are called sister chromatids.
- Centromere: the chromosomal region that holds sister chromatids together and where the kinetochore forms.




Chromosome Related Nomenclature (II)

- Kinetochores: the centromeric substructure that binds microtubules and directs chromosome movement in mitosis.
- Gene: a segment of DNA encoding a functional RNA or protein product.



Cheeseman & Desai, *Nat. Rev. Mol. Cell Biol.* 9:33, 2008


General Organization of Chromosomes





An Overview of Human Chromosomes

- Human genome

- 3.2×10^9 bp
- distributed over 24 chromosomes
- 20,000~25,000 protein coding genes



Alberts *MBoC 5e*




Organization of Human Chromosomes (I)

- Bacterial genomes make very efficient use of space. Coding sequences occupies 90% of the genome .
- For human genome, protein-coding genes occupies 1.5% of the genome.
- Human genes range in size from several hundred bp to one million bp, with an average of ~27,000 bp.

Table 4-1 Some Vital Statistics for the Human Genome

	HUMAN GENOME
DNA length	3.2×10^9 nucleotide pairs*
Number of genes	approximately 25,000
Largest gene	2.4×10^6 nucleotide pairs
Mean gene size	27,000 nucleotide pairs
Smallest number of exons per gene	1
Largest number of exons per gene	178
Mean number of exons per gene	10.4
Largest exon size	17,106 nucleotide pairs
Mean exon size	145 nucleotide pairs
Number of pseudogenes**	more than 20,000
Percentage of DNA sequence in exons (protein coding sequences)	1.5%
Percentage of DNA in other highly conserved sequences***	3.5%
Percentage of DNA in high-copy repetitive elements	approximately 50%




Organization of Human Chromosomes (II)

- Number of genes per one million bp

- 483 in budding yeast
- 197 in *C. elegans*
- 117 in fruit fly
- 7~9 in human


- Distributions of genes on chromosomes are highly variable.

- On human chromosome 9, 3~22 genes per one million bp
- One region on human chromosome 21 with 7×10^6 bp has no genes.




Organization of Human Chromosomes (III)

- Much (40~50%) of non-protein-coding DNA in the human genome is transcribed into RNA.
- Coding regions are usually unique.
- Eukaryotic genomes often contain large numbers of repetitive DNA sequences that are present in many copies.
 - Transposons
 - Satellite DNAs (e.g. centromeres)



Organization of Human Chromosomes (IV)

- Pseudogenes are DNA fragments that contain too many mutations that render a n ancestral gene inactive and nonfunctional.
- Transposons are DNA segments that can move from one location of the genome to another location.
- There are three types of transposons
 - DNA-only transposons
 - Retroviral-like retrotransposons
 - Nonretroviral-like retrotransposons




Outline

- Structural and functional organization of chromosomes
- **DNA packaging and structural organization**
- Overview of kinetochore structure and functions


DNA Packaging

- Long sequence of DNA must be stored within the geometry of a nucleus
 - Example: human chromosome 22, 48 million bp
 - Extends to length of ~1.5 cm
 - Measures 2 μm in mitosis
 - Packaging ratio on the level of 10^4 in mitosis
 - Packaging ratio ~500 in interphase
- Packaged DNA must provide for gene expression.




Packaging of DNA into Nucleosomes

- DNA is coiled around a protein core to form nucleosomes: 142 hydrogen bonds.
- ~7 folds in packaging.
- Histone H2A, H2B, H3, H4 with 147 bp DNA.
- Nucleosomes repeat at every 200 bp. So ~30 million nucleosomes in a human cell.
- Total mass of histones approximately equal to that of DNA.



Histone Organization

- N-terminal tail is subject to different forms of modification.
- H2A & H2B form a dimer through handshaking.
- H3 & H4 form a dimer in a similar fashion.
- Extensive interactions between histones and DNA





Formation of the Histone Octamer

- Histone code hypothesis:

Histone modifications determine whether particular regions of chromatin are transcribed or held in an inactive state.


- Nucleosomes are dynamic.

- Eukaryotic cells have a large variety of ATP-dependent chromatin remodeling complexes.



Second level of DNA Packaging: 30-nm Fibre

- Nucleosomes are further packaged into 30-nm fibers.
- The precise structure of the 30-nm fiber is not yet known.
- Chromatin structure beyond nucleosomes is poorly understood.





Different models of the 30-nm fiber


- ~40 folds in packaging.

Higher Order Structure of Chromosomes (I)


- Two different cases
 - Interphase chromosomes
 - Mitotic chromosomes
- Interphase chromosomes
 - Formation of loops
 - Formation of bands
- Higher order structures are also actively regulated.



Higher Order Structure of Chromosomes (II)





Model of mitotic chromosomes




The Nuclear Matrix

- The identification of nuclear matrix is controversial.
- The proposed function of nuclear matrix is to organize chromosomes compartments and chromatin loops.



Mitotic Chromosome Scaffold Protein: SMC


- SMC protein: structural maintenance of chromosome.
- Condensin: mediates condensation
- Cohesin: holds sister chromatids together; cleaved by separase.
- Hundreds of other chromosome scaffold proteins with unknown functions.




Hirano T, *Genes Dev.* 13:11-19, 1999

Structural Compartmentation of the Nucleus

- Individual chromosomes tend to concentrate within discrete territories with limited intermingling.
- Chromosomes active in transcription (euchromatin) tend to concentrate to the middle of the nucleus.
- Chromosomes inactive (heterochromatin) in transcription tend to concentrate at the periphery.



Example: Genetic vs Epigenetic Inheritance





Outline

- Structural and functional organization of chromosomes
- DNA packaging and structural organization
- **Overview of kinetochore structure and functions**

Kinetochores (I)


- Kinetochores become visible in mitosis.



Cheeseman & Desai, *Nat. Rev. Mol. Cell Biol.* 9:33, 2008

Kinetochores (II)


- Embedded in the surface of centromere.
- Kinetochores assemble in prophase; disassemble after mitosis.
- Primary functions
 - Directs chromosome movement
 - Regulating microtubule dynamics
 - Form signaling pathways to regulate cell cycle
- Fibrous corona is detected on unattached kinetochores.



Cheeseman & Desai, *Nat. Rev. Mol. Cell Biol.* 9:33, 2008

Kinetochores (III)

- Kinetochores are a very large protein assembly.
- At least 70 kinetochore-associated proteins have been identified in budding yeast.
- Its structure is likely to be fully solved in the near future.





Massachio & Salmon, *Nat. Rev. Mol. Cell Biol.* 8:379, 2007

Chromosome Organization & DNA Packaging


Transposons and Retrotransposons

- Transposons make much of human genome.
- Transposons are small discrete DNA elements that are capable of moving along DNA. Most, if not all, of them are inactive in human genome.
- Transposons move in a cut-and-paste fashion.
- Retrotransposons move via RNA intermediates.
- Retrotransposons move in a copy-and-paste fashion.



Pseudogenes

- Pseudogenes are non-functional derivatives of genes.
- Processed pseudogenes are created by reverse transcription of mature mRNA.
- Unprocessed pseudogenes are created by reverse transcription of pre-mRNA.
- Human genome contains many pseudogenes, perhaps more than functional genes.




Background Information: Epigenetics & ISH

- Epigenetics refer to changes in phenotype or gene expression caused by factors other than DNA sequence differences.
- One example: a fertilized egg cell can differentiate into many different cell types.
- In situ hybridization: the technique of using a labeled complementary DNA/RNA to specifically match with a specific DNA/RNA sequence in cells or tissues.



Background Information: Budding Yeast and Fission Yeast

- No nuclear envelope breakdown during yeast cell division.

(A) FISSION YEAST (*Schizosaccharomyces pombe*)



(B) BUDDING YEAST (*Saccharomyces cerevisiae*)



10 μ m